

Homework 1

Jean-Philippe Vert

Due January 19, 2009

1 Some kernels

Which of the following functions are positive definite:

$$\forall x, y \geq 0 \quad K_1(x, y) = \min(x, y)$$

$$\forall x, y \geq 0 \quad K_2(x, y) = \max(x, y)$$

$$\forall x, y > 0 \quad K_3(x, y) = \frac{\min(x, y)}{\max(x, y)}$$

$$\forall x, y > 0 \quad K_4(x, y) = \frac{\max(x, y)}{\min(x, y)}$$

2 Completeness of the RKHS

We want to finish the construction of the RKHS associated to a positive definite kernel K given in the course. Remember we have defined the set of functions:

$$\mathcal{H}_0 = \left\{ \sum_{i=1}^n \alpha_i K_{x_i} : n \in \mathbb{N}, \alpha_1, \dots, \alpha_n \in \mathbb{R}, x_1, \dots, x_n \in \mathcal{X} \right\}$$

and for any two functions $f, g \in \mathcal{H}_0$, given by:

$$f = \sum_{i=1}^m a_i K_{\mathbf{x}_i}, \quad g = \sum_{j=1}^n b_j K_{\mathbf{y}_j},$$

we have defined the operation:

$$\langle f, g \rangle_{\mathcal{H}_0} := \sum_{i,j} a_i b_j K(\mathbf{x}_i, \mathbf{y}_j).$$

In the course we have shown that \mathcal{H}_0 endowed with this inner product is a pre-Hilbert space. Let us now show how to finish the construction of the RKHS from \mathcal{H}_0

1. Show that any Cauchy sequence (f_n) in \mathcal{H}_0 converges pointwisely to a function $f : \mathcal{X} \rightarrow \mathbb{R}$ defined by $f(x) = \lim_{n \rightarrow +\infty} f_n(x)$.

2. Show that any Cauchy sequence $(f_n)_{n \in \mathbb{N}}$ in \mathcal{H}_0 which converges pointwisely to 0 satisfies:

$$\lim_{n \rightarrow +\infty} \|f_n\|_{\mathcal{H}_0} = 0.$$

3. Let $\mathcal{H} \subset \mathbb{R}^{\mathcal{X}}$ be the set of functions $f : \mathcal{X} \rightarrow \mathbb{R}$ which are pointwise limits of Cauchy sequences in \mathcal{H}_0 , i.e., if (f_n) is a Cauchy sequence in \mathcal{H}_0 , then $f(x) = \lim_{n \rightarrow +\infty} f_n(x)$. Show that $\mathcal{H}_0 \subset \mathcal{H}$.

4. If (f_n) and (g_n) are two Cauchy sequences in \mathcal{H}_0 , which converge pointwisely to two functions f and $g \in \mathcal{H}$, show that the inner product $\langle f_n, g_n \rangle_{\mathcal{H}_0}$ converges to a number which only depends on f and g . This allows us to define formally the operation:

$$\langle f, g \rangle_{\mathcal{H}} = \lim_{n \rightarrow +\infty} \langle f_n, g_n \rangle_{\mathcal{H}_0}.$$

5. Show that $\langle \cdot, \cdot \rangle_{\mathcal{H}}$ is an inner product on \mathcal{H} .

6. Show that \mathcal{H}_0 is dense in \mathcal{H} (with respect to the metric defined by the inner product $\langle \cdot, \cdot \rangle_{\mathcal{H}}$)

7. Show that \mathcal{H} is complete.

8. Show that \mathcal{H} is a RKHS whose reproducing kernel is K .

3 Unicity of the RKHS (Bonus)

Prove that if $K : \mathcal{X} \times \mathcal{X}$ is a positive definite function, then it is the r.k. of a unique RKHS. (Hint: consider the linear space spanned by the functions $K_x : t \mapsto K(x, t)$, and use the fact that a linear subspace \mathcal{F} of a Hilbert space \mathcal{H} is dense in \mathcal{H} if and only 0 is the only vector orthogonal to all vectors in \mathcal{F})