

MicroRNA-Cancer Connection: The Beginning of a New Tale

George Adrian Calin and Carlo Maria Croce

Department of Molecular Virology, Immunology, and Medical Genetics and Comprehensive Cancer Center, Ohio State University, Columbus, Ohio

Abstract

Cancer initiation and progression can involve microRNAs (miRNA), which are small noncoding RNAs that can regulate gene expression. Their expression profiles can be used for the classification, diagnosis, and prognosis of human malignancies. Loss or amplification of miRNA genes has been reported in a variety of cancers, and altered patterns of miRNA expression may affect cell cycle and survival programs. Germ-line and somatic mutations in miRNAs or polymorphisms in the mRNAs targeted by miRNAs may also contribute to cancer predisposition and progression. We propose that alterations in miRNA genes play a critical role in the pathophysiology of many, perhaps all, human cancers. (Cancer Res 2006; 66(15): 7390-4)

Cancer-Specific MicroRNA Fingerprints

Cancer is a very complex genetic disease characterized by alterations in genes encoding oncogenic and tumor-suppressor proteins [protein coding genes (PCG)]. First described in *C. elegans* more than a decade ago (1), >3,000 members of a new class of small noncoding RNAs, named microRNAs (miRNAs; ref. 2), have been identified in the last 5 years in vertebrates, flies, worms, and plants, and even in viruses. Functionally, it was shown that miRNAs reduce the levels of many of their target transcripts as well as the amount of protein encoded by these transcripts (3). For several miRNAs, the participation in essential biological processes has been proved, such as cell proliferation control (*miR-125b* and *let-7*), hematopoietic B-cell lineage fate (*miR-181*), B-cell survival (*miR-15a* and *miR-16-1*), brain patterning (*miR-430*), pancreatic cell insulin secretion (*miR-375*), and adipocyte development (*miR-143*; for reviews, see ref. 4).

After the identification of two clustered miRNAs as the targets of homozygous and heterozygous deletions and translocations at 13q14.3 in human B-cell chronic lymphocytic leukemias (B-CLL; ref. 5), the question to be answered was how general is the involvement of miRNAs in human cancers. The development of miRNA microarrays was a necessary step for the high-throughput miRNA fingerprint investigation in normal and cancer cells (6). Other technologies, including macroarrays (7), bead-based flow cytometric miRNA expression (8), and quantitative reverse transcription-PCR (9), are now available. What we learned from such expression studies is reshaping the landscape of cancer genomics (Table 1 and included references).

Cancer-specific miRNA fingerprints were identified in every type of analyzed cancer, including B-CLL (10), breast carcinoma (11), primary glioblastoma (12), hepatocellular carcinoma (13), papillary thyroid carcinoma (14), lung cancer (15–17), gastric carcinoma,

colon carcinoma (18), and endocrine pancreatic tumors (17). Not only the spectrum of miRNAs expressed in malignant cells is significantly different from that of normal counterpart cells but also miRNA expression profiles better classify poorly differentiated tumors as compared with the mRNA (EST)-based classifier (8). Commonly deregulated miRNAs in different types of solid cancers predict their involvement in fundamental pathways and their interaction with important cancer-specific PCGs (17). Furthermore, such abnormal expression was found not only in malignant cells but also in premalignant stages, such as colon adenomas where *miR-143* and *miR-145* expression is reduced (18) or in pituitary adenomas, a type of benign tumors displaying deletions at 13q14.3 and reduced expression of *miR-16-1* and *miR-15a* (19). The finding that essentially all indolent CLLs have lost *miR-15a/miR-16-1* expression suggests that this event is the initiating event in the pathogenesis of the indolent form of CLL (5, 10, 20). Furthermore, it was shown that *miR-221*, highly overexpressed in papillary thyroid tumors, is also overexpressed in normal thyroid tissue adjacent to tumors but not in normal thyroid tissues from individuals without clinical thyroid disease (14). Therefore, it seems likely that, in some cases, the cancer-specific miRNA fingerprints represent events involved in the initiation of the malignant process.

What are the causes of the widespread miRNA misexpression in cancers? Although not clearly understood, the origins of such abnormalities seem to be multiple. Many miRNAs reside in genomic regions involved in cancer, including minimal regions of loss of heterozygosity (LOH), minimal amplicons, or breakpoint cluster regions (21). As shown in Table 1, the overexpressed oncogenic miRNAs are located in amplified regions and the down-regulated suppressor miRNAs in deleted regions in cancers. The proof that chromosomal rearrangements are causal includes the early report of a masked t(8;17) translocation that resulted in an aggressive B-cell leukemia by overexpressing *c-myc* oncogene by an unknown mechanism at the moment of identification (22). It was shown later that *miR-142* is located at the chromosome 17 breakpoint and that *c-Myc* was rearranged under the control of the promoter of *miR-142* with consequent overexpression. In a precursor B-cell acute lymphoblastic leukemia, an insertion of *miR-125b-1* into a rearranged immunoglobulin heavy chain locus was described, suggesting an early involvement in leukemogenesis (23).

Mutations in MiRNAs: A Way to Predispose to Cancer?

In spite of decades of research, the molecular basis for the major fraction of familial cancers is unknown. CLL represents one of the main examples in this regard: a significant portion (10-20%) of patients have a family history of CLL or other hematologic or solid cancers whereas no clear culprit could be found by scanning PCGs (20, 24). Screening the human miRNoma for sequence abnormalities located either in the pre-miRNA or in pri-miRNA, a higher frequency of germ-line or somatic mutations (about 15%), as expected by the small size of miRNA genes was found (25).

Requests for reprints: Carlo M. Croce, Comprehensive Cancer Center, Ohio State University, Room 385K, Wiseman Hall, 400 12th Avenue, Columbus, OH 43210. Phone: 614-292-4354; Fax: 614-292-3312; E-mail: Carlo.Croce@osumc.edu.

©2006 American Association for Cancer Research.
doi:10.1158/0008-5472.CAN-06-0800

Table 1. Facts about tumor suppressor and oncogenic miRNAs

| MiRNA | Location | Putative function | CAGR location* | Cancer abnormalities/description | Reference |
|-----------------------------|---|-------------------|---|--|-----------|
| <i>miR-16-1-15a</i> cluster | 13q14.3, intron 4 of <i>DLEU2</i> | Suppressor miRNAs | LOH in CLL and prostate cancer | Deleted and down-regulated in the majority of B-CLLs | (5) |
| | | | | Reduced expression in the majority of DLBCLs | (31) |
| | | | | Down-regulation in pituitary adenomas | (19) |
| | | | | Reduced expression associated with good prognosis in B-CLL | (25) |
| | | | | Germ-line mutations in the primary transcript <i>miR-16-1/15a</i> in B-CLLs | (25) |
| | | | | Exogenous restoration in leukemia cells of <i>miR-16/15</i> induces apoptosis by directly targeting BCL2 | (36) |
| <i>miR-145</i> | 5q32, intergenic | Suppressor miRNA | LOH in MDS (5q- syndrome) | Reduced accumulation in colon adenomas and carcinomas | (18) |
| | | | | Reduced expression in breast cancers | (11) |
| <i>let-7</i> family | various | Suppressor miRNAs | LOH in lung cancers | Reduced expression associated with shortened postoperative survival | (15) |
| | | | | <i>let-7a-1</i> expression correlates with poor survival of lung cancer patients | (16) |
| | | | | <i>let-7</i> regulates RAS oncogene expression in lung tumors | (33) |
| <i>miR-155</i> | 21q21.3, exon 3 of noncoding RNA <i>BIC</i> | Oncogenic miRNA | not reported | High expression of precursor <i>miR-155/BIC</i> in pediatric BL, but lack of <i>BIC</i> and <i>miR-155</i> expression in adult BL | (38, 41) |
| | | | | <i>miR-155</i> overexpressed in B-cell lymphomas, significantly higher levels in DLCL with poor prognosis (activated B-cell phenotype) | (31, 39) |
| | | | | Increased expression of <i>miR-155</i> in Epstein-Barr transformed lymphoblastoid cell lines | (9) |
| | | | | High expression of both <i>BIC</i> and <i>miR-155</i> in Hodgkin, primary mediastinal and DLBCL lymphomas | (32) |
| | | | | Overexpression in breast cancers | (11) |
| | | | | <i>miR-155</i> overexpression correlates with poor survival in lung cancers | (16) |
| <i>miR-17-92</i> cluster | 13q31.3, intron 3 <i>C13orf25</i> | Oncogenic miRNA | AMPLIF in follicular lymphomas | Target of genomic amplification in malignant lymphomas | (27, 28) |
| | | | | Overexpressed in lung cancers; the miRNA cluster, but not the host <i>C13orf25</i> gene, enhances cell proliferation. | (40) |
| | | | | Primary transcripts overexpressed in lymphomas, but not in colorectal carcinomas; enforced expression acted with the c-Myc expression to accelerate tumor development in mouse B-cell lymphomas. | (29) |
| | | | | Negative regulatory feed-back loop c-Myc/miR-17-5p-miR-20a/E2F1 | (37) |
| <i>miR-21</i> | 7q23.2, 3'UTR VMP1 | Suppressor miRNAs | AMPLIF in neuroblastoma and breast cancer | Elevated levels in glioblastoma primary tumors and cell lines; increased apoptotic cell death after <i>miR-21</i> knockdown in glioblastoma cells | (7) |
| | | Oncogenic miRNA | | | |

Abbreviations: B-CLL, B-cell chronic lymphocytic leukemia; *BIC*, noncoding RNA gene; BL, Burkitt lymphoma; DLBCL, diffuse large B-cell lymphoma; *DLEU2*, noncoding RNA gene; VMP1, vacuole membrane protein 1.

*CAGR, cancer-associated genomic regions (as in ref. 21).

Furthermore, a germ-line mutation in the *pri-miR-16-1/15a* precursor in a patient with familial CLL and breast cancer in first-degree members of family suggests a possible predisposing effect. The roles of mutations in miRNAs have still to be elucidated,

and tumor-specific *pri-miRNA* sequence abnormalities seem to be a more widespread phenomenon in tumorigenesis since mutations near the clusters *miR-17-92* on chromosome 13 and *miR-106-92* on chromosome X were described in a mouse model (26). It was

shown that the cluster *miR-17-92* is amplified in human lymphomas (27, 28) and accelerates *c-Myc*-induced tumorigenesis in a mouse model of B-cell lymphoma (29), suggesting a pathogenetic role of such mutations.

As the thermodynamics of RNA-RNA binding plays essential roles in the miRNA interaction with the target mRNA, it is supposed that sequence variations influencing this interaction will be identified in cancers. Thyroid cancers in which the up-regulation of *miR-221*, *miR-222*, and *miR-146* was the strongest showed dramatic loss of *KIT* oncogene and, in half of the cases, the down-regulation was associated with germ-line single-nucleotide polymorphisms in the two recognition sites in *KIT* for these three miRNAs (14). It has to be noted that thyroid papillary carcinoma is a type of cancer with high familiarity without known genetic bases. As the 3' untranslated region (UTR) of PCGs was scarcely screened for mutations/polymorphisms, it is possible that the extent of such abnormalities might be much larger than initially thought. Further strengthening possible roles of polymorphisms in altering the function of miRNAs, a study in Japanese normal subjects screened for single-nucleotide polymorphisms in the genomic regions corresponding to 173 precursor miRNAs found a polymorphism in the mature *miR-30c-2* sequence that may alter target selection and exert biological effects (30). Making the story more intriguing, this miRNA is a member of a common expression signature characterizing several solid cancers (17). Putting all these data together, it is tempting to propose that germ-line mutations or polymorphisms in miRNA genes or interacting sequences in target mRNA might represent a newly described mechanism of cancer predisposition. Further identification of sequence or expression variations in miRNAs in a large series of familial cancer patients is needed to clearly prove this hypothesis.

MiRNAs: From the Scientist Bench to the Patient Bedside

It is well known that PCGs with important cancer connections are used also as diagnosis markers and therapy targets. If the miRNAs are active players in human oncogenesis, then they will have an effect on the diagnosis and prognosis of cancer (Table 1). In fact, evidence that miRNAs represent new diagnostic and prognostic factors in human cancers is rapidly accumulating. In B-CLL, a unique miRNA signature is associated with prognostic factors and with the time from diagnosis to initiation of therapy (25). In diffuse large B-cell lymphoma, independent studies revealed that significantly higher levels of *miR-155* occur in cases with poorer prognosis (an activated B-cell phenotype) than in those with the germinal center phenotype (31, 32). Expression of members of *let-7* family correlates with postoperative survival in lung cancer, the group of patients with reduced expression showing significantly shorter survival after potentially curative resection (15). In lung adenocarcinomas, high *miR-155* and low *let-7a-2* expression correlates with poor survival (16). In breast carcinomas, miRNA expression was correlated with specific biopathologic features, such as estrogen and progesterone receptor expression (the members of *miR-30* family), or tumor stage (*miR-213* and *miR-203*; ref. 11). Expression of three genes, *miR-92*, *miR-20*, and *miR-18*, was inversely correlated with the degree of hepatocellular carcinoma differentiation (13). Such results strongly suggest that quantification of miRNAs may be diagnostically useful.

To understand the possible role of miRNAs as putative therapeutic agents, we have to elucidate the consequences of the wide-

spread miRNA dysregulation in cancer cells. In lung cancers, activation of *RAS* genes by point mutations, identified more than two decades ago, may represent an early event in some tumors. *RAS* protein is significantly higher in lung tumors than in normal lung tissue whereas *let-7* expression is lower in lung cancer cells. This correlation led to the identification of a direct regulation of *RAS* by the *let-7* miRNA family (33). Exogenous delivery of *let-7* to the lung might either prevent the formation of lung tumors (from pre-malignant lesions) or shrink tumors with activating *RAS* mutations (34).

MiRNAs are natural antisense interactors with players in the eukaryotic survival and cell cycle programs. The overexpression of antiapoptotic protein *BCL2* is an important genetic event in human tumorigenesis, including follicular lymphoma, lung cancer, and B-CLL. The mechanism of this activation, except in all cases of follicular lymphomas where a translocation t(14;18) is responsible (35), was unknown. Loss of *miR-15a/miR-16-1* in CLL results in *BCL2* overexpression and restoration of *miR-15/miR-16* in leukemia cells induces apoptosis by directly interacting with *BCL2* mRNA (36). These results are encouraging in the light of new promising results on the therapeutic potential of antisense *BCL2*.

The oncogene *c-myc* encodes a transcription factor that regulates, via several targets including E2F1 transcription factor, cell proliferation and survival. A feedback regulatory loop in which *MYC* directly binds and activates the transcription of the cluster *miR-17-92* that consequently negatively regulates E2F1 by direct interaction, while *c-Myc* is directly inducing expression of the E2F1 that in turn induces *c-Myc*, was recently described (37). This fine molecular dissection of an important cellular pathway has cancer implications, as it was shown that *c-myc* and *miR-17-92* cooperate and such cooperation accelerates B-cell tumorigenesis in a mouse lymphoma model (29). Such results offer a rationale basis for targeted therapy (e.g., by using antisense miRNAs against the clustered miRNAs) that will overload the regulatory loop, with the acceleration of the *MYC*-E2F1 feedback and consequent cell death.

The "MiRNA Cascade": A Model of MiRNA Involvement in Human Cancers

MiRNAs are contributors to oncogenesis, functioning as tumor suppressors (as is the case of *miR-15a* and *miR-16-1*) or as oncogenes (as is the case of *miR-155* or *miR-17-92* cluster; Fig. 1; refs. 21, 34). The classic tumorigenesis model postulates alterations in protein-coding oncogenes and tumor suppressor genes. Relatively minor variations in the levels of expression of a miRNA or mutations that affect moderately the conformation of miRNA:mRNA pairing could have important consequences for the cell. The explanation is provided by the large number of targets of each miRNA and the relatively large number of altered miRNAs, making very likely that two or more PCGs from different molecular pathways/interacting pathways are disturbed. The down-regulation of the suppressor *miR-15a/miR-16-1* induces overexpression of *BCL2* and possibly other genes that may be important for tumorigenesis, whereas the overexpression of oncogenic *miR-17-92* cooperates with *c-myc* in stimulating proliferation. Therefore, the miRNAs may act "in cascade" over several cancer-specific PCGs, which in turn could influence the transcription or function of several other PCGs and noncoding RNAs including miRNAs. If miRNA alterations occur in somatic cells, they could initiate or contribute to tumorigenesis, whereas, if present in the germ-line, could represent cancer-predisposing events. A paradigm for this model is human B-CLL, in which *miR-15a* and *miR-16-1* are located




Figure 1. miRNA activation and inactivation events and cooperation with protein coding genes in human tumorigenesis. The abnormalities found to influence the activity of miRNAs are the same as those described to target PCGs, including chromosomal rearrangements, genomic amplifications or deletions, and mutations. In a specific tumor, both abnormalities in PCGs and miRNAs can be identified. Inactivation of tumor suppressor PCGs and activation of oncogenic miRNAs have the same molecular consequences: reduced levels of proteins blocking proliferation and activating apoptosis. By contrast, activation of oncogenic PCGs and inactivation of suppressor miRNAs are followed by accumulation of proteins that stimulate proliferation and decrease apoptosis. For example, effects of t(14;18)(q32;q21) or del13q14.3 in leukemic cells are the same: overexpression of the antiapoptotic *BCL2* protein, in the former case by juxtaposition of oncogene *BCL2* to immunoglobulin enhancers and in the latter by down-regulation of suppressor *miR-16-1* and *miR-15a*, which negatively regulate *BCL2* production. Triangles, promoter regions; circles and rectangles, miRNA and PCG structural genes.

in the most frequently deleted genomic region, are down-regulated in the majority of cases, harbor mutations in familial cases, and induce apoptosis in a leukemia model by targeting the overexpressed antiapoptotic *BCL2* gene. As the puzzle of noncoding RNA involvement in cancer is just starting to be assembled, certainly further unexpected pieces will be identified in the near future.

Acknowledgments

Received 3/1/2006; revised 4/21/2006; accepted 5/4/2006.

Grant support: Program Project Grants from the National Cancer Institute (C.M. Croce) and Kimmel Foundation Scholar award and CLL Global Research Foundation (G.A. Calin).

We apologize to our many colleagues whose works were not cited due to space limitations.

References

- Lee RC, Feinbaum RL, Ambros V. The *C. elegans* heterochronic gene *lin-4* encodes small RNAs with antisense complementarity to *lin-14*. *Cell* 1993;75:843–54.
- Ambros V. MicroRNA pathways in flies and worms: growth, death, fat, stress, and timing. *Cell* 2003;113:673–6.
- Lim LP, Lau NC, Garrett-Engle P, et al. Microarray analysis shows that some microRNAs down-regulate large numbers of target mRNAs. *Nature* 2005;433:769–73.
- Harfe BD. MicroRNAs in vertebrate development. *Curr Opin Genet Dev* 2005;15:410–5.
- Calin GA, Dumitru CD, Shimizu M, et al. Frequent deletions and down-regulation of micro-RNA genes *miR15* and *miR16* at 13q14 in chronic lymphocytic leukemia. *Proc Natl Acad Sci U S A* 2002;99:15524–9.
- Liu C-G, Calin GA, Meloon B, et al. An oligonucleotide microchip for genome-wide miRNA profiling in human and mouse tissues. *Proc Natl Acad Sci U S A* 2004;101:9740–4.
- Chan JA, Krichevsky AM, Kosik KS. MicroRNA-21 is an antiapoptotic factor in human glioblastoma cells. *Cancer Res* 2005;65:6029–33.
- Lu J, Getz G, Miska EA, et al. MicroRNA expression profiles classify human cancers. *Nature* 2005;435:834–8.
- Jiang J, Lee EJ, Schmittgen TD. Increased expression of microRNA-155 in Epstein-Barr virus transformed lymphoblastoid cell lines. *Genes Chromosomes Cancer* 2005;45:103–6.
- Calin GA, Liu CG, Sevignani C, et al. MicroRNA profiling reveals distinct signatures in B cell chronic lymphocytic leukemias. *Proc Natl Acad Sci U S A* 2004;101:11755–60.
- Iorio MV, Ferracin M, Liu CG, et al. microRNA gene expression deregulation in human breast cancer. *Cancer Res* 2005;65:7065–70.
- Ciafre SA, Galardi S, Mangiola A, et al. Extensive modulation of a set of microRNAs in primary glioblastoma. *Biochem Biophys Res Commun* 2005;334:1351–8.
- Murakami Y, Yasuda T, Saigo K, et al. Comprehensive analysis of microRNA expression patterns in hepatocellular carcinoma and non-tumorous tissues. *Oncogene* 2006;25:2537–45.
- He H, Jazdzewski K, Li W, et al. The role of microRNA genes in papillary thyroid carcinoma. *Proc Natl Acad Sci U S A* 2005;102:19075–80.
- Takamizawa J, Konishi H, Yanagisawa K, et al. Reduced expression of the let-7 microRNAs in human

- lung cancers in association with shortened postoperative survival. *Cancer Res* 2004;64:3753–6.
16. Yanaihara N, Caplen N, Bowman E, et al. microRNA signature in lung cancer diagnosis and prognosis. *Cancer Cell* 2006;9:189–98.
17. Volinia S, Calin GA, Liu C-G, et al. A microRNA expression signature of human solid tumors define cancer gene targets. *Proc Natl Acad Sci U S A* 2006; 103:2257–61.
18. Michael MZ, O'Connor SM, van Holst Pellekaan NG, Young GP, James RJ. Reduced accumulation of specific microRNAs in colorectal neoplasia. *Mol Cancer Res* 2003;1:882–91.
19. Bottoni A, Piccin D, Tagliati F, Luchin A, Zatelli MC, degli Uberti EC. miR-15a and miR-16-1 down-regulation in pituitary adenomas. *J Cell Physiol* 2005; 204:280–5.
20. Calin GA, Trapasso F, Shimizu M, et al. Familial cancer associated with a polymorphism in ARLTS1. *New Engl J Med* 2005;352:1667–76.
21. Calin GA, Sevignani C, Dumitru CD, et al. Human microRNA genes are frequently located at fragile sites and genomic regions involved in cancers. *Proc Natl Acad Sci U S A* 2004;101:2999–3004.
22. Gauwerky CE, Huebner K, Isobe M, Nowell PC, Croce CM. Activation of MYC in a masked t(8;17) translocation results in an aggressive B-cell leukemia. *Proc Natl Acad Sci U S A* 1989;86:8867–71.
23. Sonoki T, Iwana E, Mitsuya H, Asou N. Insertion of microRNA-125b-1, a human homologue of lin-4, into a rearranged immunoglobulin heavy chain gene locus in a patient with precursor B-cell acute lymphoblastic leukemia. *Leukemia* 2005;19:2009–10.
24. Catovsky D. Definition and diagnosis of sporadic and familial chronic lymphocytic leukemia. *Hematol Oncol Clin North Am* 2004;8:783–94.
25. Calin GA, Ferracin M, Cimmino A, et al. A unique microRNA signature associated with prognostic factors and disease progression in B cell chronic lymphocytic leukemia. *N Engl J Med* 2005;352:1667–76.
26. Kool J, Lagcher W, Uren AG, et al. Mutation of miRNA genes in high-throughput *in vivo* screens for genes involved in tumorigenesis. In: *Keystone Symposia: RNAi and related pathways*. Vancouver, British Columbia; 2006.
27. Ota A, Tagawa H, Karnan S, et al. Identification and characterization of a novel gene, C13orf25, as a target for 13q31-32 amplification in malignant lymphoma. *Cancer Res* 2004;64:3087–95.
28. Tagawa H, Seto M. A microRNA cluster as a target of genomic amplification in malignant lymphoma. *Leukemia* 2005;19:2013–6.
29. He L, Thomson JM, Hemann MT, et al. A microRNA polycistron as a potential human oncogene. *Nature* 2005;435:828–33.
30. Iway N, Naraba H. Polymorphisms in human pre-miRNAs. *Biochem Biophys Res Commun* 2005;331: 1439–44.
31. Eis PS, Tam W, Sun L, et al. Accumulation of miR-155 and BIC RNA in human B cell lymphomas. *Proc Natl Acad Sci U S A* 2005;102:3627–32.
32. Kluiver J, Poppema S, de Jong D, et al. BIC and miR-155 are highly expressed in Hodgkin, primary mediastinal and diffuse large B cell lymphomas. *J Pathol* 2005; 207:243–9.
33. Johnson SM, Grosshans H, Shingara J, et al. J. RAS is regulated by the let-7 microRNA family. *Cell* 2005;120: 635–47.
34. Esquela-Kerscher A, Slack FJ. Oncomirs—microRNAs with a role in cancer. *Nat Rev Cancer* 2006;6:259–69.
35. Tsujimoto Y, Finger LR, Yunis J, Nowell PC, Croce CM. Cloning of the chromosome breakpoint of neoplastic B cells with the t(14;18) chromosome translocation. *Science* 1984;226:1097–9.
36. Cimmino A, Calin GA, Fabbri M, et al. miR-15 and miR-16 induce apoptosis by targeting BCL2. *Proc Natl Acad Sci U S A* 2005;102:13944–9.
37. O'Donnell KA, Wentzel EA, Zeller KI, Dang CV, Mendell JT. c-Myc-regulated microRNAs modulate E2F1 expression. *Nature* 2005;435:839–43.
38. Metzler M, Wilda M, Busch K, Viehmann S, Borkhardt A. High expression of precursor microRNA-155/BIC RNA in children with Burkitt lymphoma. *Genes Chromosomes Cancer* 2004;39:167–9.
39. Tam W, Dahlberg JE. miR-155/BIC as an oncogenic microRNA. *Genes Chromosomes Cancer* 2006;45:211–2.
40. Hayashita Y, Osada H, Tatematsu Y, et al. A polycistronic microRNA cluster, miR-17-92, is overexpressed in human lung cancers and enhances cell proliferation. *Cancer Res* 2005;65:9628–32.
41. Kluiver J, Haralambieva E, de Jong D, et al. Lack of BIC and microRNA miR-155 expression in primary cases of Burkitt lymphoma. *Genes Chromosomes Cancer* 2006;45:147–53.